

**Priority Health Medicare prior authorization form**

Fax completed form to: 877.974.4411 toll free, or 616.942.8206

This form applies to:  Medicare Part B  Medicare Part D  
 This request is:  Expedited request  Standard request

Your request will be expedited if you haven't gotten the prescription and Priority Health Medicare determines, or your prescriber tells us, that your life or health may be at risk by waiting.

**Testosterone topical**

**Member**

Last Name: \_\_\_\_\_ First Name: \_\_\_\_\_  
 ID #: \_\_\_\_\_ DOB: \_\_\_\_\_ Gender: \_\_\_\_\_  
 Primary Care Physician: \_\_\_\_\_  
 Requesting Provider: \_\_\_\_\_ Prov. Phone: \_\_\_\_\_ Prov. Fax: \_\_\_\_\_  
 Provider Address: \_\_\_\_\_  
 Provider NPI: \_\_\_\_\_ Contact Name: \_\_\_\_\_  
 Provider Signature: \_\_\_\_\_ Date: \_\_\_\_\_

**Drug information**

New request  Continuation request

Drug product:  Testosterone 10mg (2%) gel **Start date** (or date of next dose): \_\_\_\_\_  
 Testosterone 12.5 mg (1%) gel **Date of last dose** (if applicable): \_\_\_\_\_  
 Testosterone 25 mg/2.5 gm (1%) gel **Dosing frequency:** \_\_\_\_\_  
 Testosterone 50 mg/5 gm (1%) gel  
 Testosterone 30 mg topical solution

**Prior authorization criteria**

The following requirements need to be met before this drug is covered by Priority Health Medicare. These requirements have been approved by the Centers for Medicare and Medicaid Services (CMS), but you may ask us for an exception if you believe one or more of these requirements should be waived.

**Before this drug is covered, the patient must meet all of the following requirements:**

1. Must have a diagnosis of hypogonadism or another medically-accepted indication\*
2. Must be male
3. Must have a serum total testosterone level less than 300 ng/dL on more than one occasion in the past year
4. Must have clinical signs and symptoms consistent with androgen deficiency
5. Must be screened for prostate cancer before starting therapy and routinely while on therapy (if appropriate):
  - Required for men over age 50
  - Required for men over age 40 who are African-American or have a family history of prostate cancer
6. Must have a documented trial and failure to a generic injectable testosterone product\*\*
  - Failure is an intolerance or an inability to improve symptoms or testosterone levels after 2 months of therapy

\*\*Note: This requirement only applies to members who are enrolled in a MAPD (Medicare Advantage Prescription Drug) plan.

**Additional information**

**Note:** When criteria are met, duration of approval is 1 year

**Medically-accepted indication\***

This drug is only covered under Medicare Part D when it is used for a medically accepted indication. A medically accepted indication is a use of the drug that is *either*:

- approved by the Food and Drug Administration. (That is, the Food and Drug Administration has approved the drug for the diagnosis or condition for which it is being prescribed.)
- — or — supported by certain reference books. (These reference books are the American Hospital Formulary Service Drug Information and the DRUGDEX Information System)

**Priority Health Precertification Documentation**

**A. What is the patient's diagnosis?**

- Hypogonadism  
 Other – the patient's condition is: \_\_\_\_\_

**Rationale for Other use:** \_\_\_\_\_

**B. Has the patient had more than one serum total testosterone level < 300 ng/dL in the past year?**

- Yes

Lab Date	Result	Interpretation		
_____	_____ ng/dL	<input type="checkbox"/> high	<input type="checkbox"/> low	<input type="checkbox"/> within normal limits
_____	_____ ng/dL	<input type="checkbox"/> high	<input type="checkbox"/> low	<input type="checkbox"/> within normal limits

- No. **Are you requesting an exception to the criteria?**  
 Yes. **Rationale for exception:** \_\_\_\_\_  
 No

**C. Does the patient have clinical signs and symptoms of hypogonadism?**

- Yes
- |   |   |
|---|---|
| <input type="checkbox"/> Decrease in energy | <input type="checkbox"/> Decrease in muscle mass  |
| <input type="checkbox"/> Fatigue            | <input type="checkbox"/> Difficulty concentrating |
| <input type="checkbox"/> Gynecomastia       | <input type="checkbox"/> Hot flashes              |
| <input type="checkbox"/> Other: _____       |   |

- No. **Are you requesting an exception to the criteria?**  
 Yes. **Rationale for exception:** \_\_\_\_\_  
 No

**D. If appropriate based on risk factors and age, has the patient been screened for prostate cancer?**

- Yes  
 No. **Are you requesting an exception to the criteria?**  
 Yes. **Rationale for exception:** \_\_\_\_\_  
 No

**E. Has the patient tried and failed a generic injectable testosterone product?\***

- Yes. Check all that apply:  
 Testosterone cypionate  
 Testosterone enanthate  
 No. **Are you requesting an exception to the criteria?**  
 Yes. **Rationale for exception:** \_\_\_\_\_  
 No

**F. What was the reason for failure to the generic injectable testosterone product?\***

- Intolerance – *Reaction:* \_\_\_\_\_
- Inability to improve symptoms or testosterone levels after 2 months of therapy

\*\*Note: This requirement only applies to members who are enrolled in a MAPD (Medicare Advantage Prescription Drug) plan.

---

**Priority Health Medicare Exception Request** (*exceptions to the above criteria*)

**Do you believe one or more of the prior authorization requirements should be waived?**  Yes  No

If yes, you must provide a statement explaining the medical reason why the exception should be approved.

**Would testosterone topical gel or solution likely be the most effective option for this patient?**

Yes  No

If yes, please explain why: \_\_\_\_\_

---

---

**If the patient is currently using testosterone topical gel or solution, would changing the patient's current regimen likely result in adverse effects for the patient?**

Yes  No

If yes, please explain: \_\_\_\_\_

---

---