PriorityHealth

Osteoporosis Management in Women Who Had a Fracture (OMW)

Women who suffer a fracture are at increased risk of additional fractures and more likely to have osteoporosis. The "Osteoporosis Management in Women Who Had a Fracture" (OMW) Star Ratings measure evaluates women 67-85 years of age who had a fracture* and had either:

- A bone mineral density (BMD) test, or
- Received a prescription to treat osteoporosis within six months of ER or inpatient discharge for a fracture

Product lines	Quality programs affected	Collection and reporting method
Medicare	CMS Star Ratings	Administrative • Claim data • Pharmacy data

How to comply with this measure

To comply with this measure, a patient must have a BMD test or be prescribed at least one of the following osteoporosis medications within 180 days of their discharge for a fracture:

Drug category	Medications	
Bisphosphonates	• Alendronate • Alendronate-cholecalciferol • Ibandronate	• Risedronate • Zoledronic acid
Other Agents	• Abaloparatide • Denosumab • Raloxifene	• Romosozumab • Teriparatide

Billing codes

Description	Code type	Codes**
Bone Mineral Density Tests	CPT	76977, 77078, 77080, 77081, 77085, 77086
Osteoporosis Medications	HCPCS	J0897, J1740, J3110, J3111, J3489

Additional information

Intake period	• July 1st of the year prior to June 30th of the measurement year
Test, service or procedure to close care opportunity	• BMD test Submit medical record information to Priority Health HEDIS department via: **Email:** *HEDIS@PriorityHealth.com* **Fax:** 616-975-8897 **Mail:** HEDIS at 1239 E Beltline, NE Mail Stop 1280, Grand Rapids, MI 49525
Medical record detail (including but not limited to)	• Bone density reports, dated with results • Medication list

SUPPLEMENTAL DATA ACCEPTED



Tips and best practices

To help close this care opportunity, we recommend you:

- Review the faxed DXA Bone Density Scan report which identifies patients who have had a recent fracture
- Have your patient complete a BMD test within six months of a fracture
- Prescribe osteoporosis medications within six months of fracture when clinically appropriate if patient is unable or unwilling to have the BMD test
- Follow up with patients who have overdue BMD referral orders and help resolve barriers to getting tested
- If the patient has an inpatient stay, request that the hospital perform a BMD test prior to discharge
- Educate member on safety and fall prevention

* Fractures of finger, toe, face and skull are not included in this measure
**Billing codes listed are for HEDIS compliance and are subject to plan coverage and contracted fee schedule

©2021 Priority Health 10761N 09/21